

CALGB-40101:

CYCLOPHOSPHAMIDE AND DOXORUBICIN (CA X 4 CYCLES) VERSUS PACLITAXEL (4 CYCLES) AS ADJUVANT THERAPY FOR BREAST CANCER IN WOMEN WITH 0-3 POSITIVE AXILLARY LYMPH NODES: A PHASE III RANDOMIZED STUDY

ClinicalTrials.gov Identifier: NCT00041119

Study Background

Trial Design: This study was conducted using a phase III factorial design. The factors are chemotherapy agent and treatment length. Cyclophosphamide plus doxorubicin (CA) is the standard agent; the experimental agent is paclitaxel (T). The standard length is shorter (4 cycles) treatment duration; the experimental length is longer (6 cycles) duration. The trial is designed to test (1) the equivalence of the experimental agent T with the standard agent combination CA; and (2) the superiority of longer versus shorter treatment duration. Patients will be randomized with equal probability to one of the four possible treatment arms.

Objectives: Primary:

1. To determine the equivalence of paclitaxel given every two weeks with CA given every two weeks as adjuvant therapy for women with 0-3 positive axillary lymph nodes, for disease-free survival.
2. To determine if longer therapy, 12 weeks, is superior to shorter therapy, 8 weeks, of either CA or paclitaxel for disease-free survival for women with primary breast cancer with 0-3 positive axillary lymph nodes.

Secondary:

- To determine the equivalence of paclitaxel given every two weeks with CA given every two weeks, and the potential superiority of longer vs. shorter therapy, in relation to overall survival, local control (regardless of metastatic status) and time to distant metastases (regardless of local recurrence status).
- To compare toxicities of short and long course CA and paclitaxel as adjuvant therapy for women with 0-3 positive axillary lymph node breast cancer.
- To determine the effect of long and short course CA and paclitaxel on the induction of menopause for pre-menopausal patients.

Stratification

Factors:

- Menopausal status (pre vs. post)
- ER/PgR status (either positive or unknown vs both negative)
- HER-2 status (positive, negative or unknown)

Study

06/27/2002 Activation Date

History:

07/30/2010 Close Date

Publication Information

Analysis Type: Primary Endpoint #1: To determine the equivalence of paclitaxel given every two weeks with CA given every two weeks as adjuvant therapy for women with 0-3 positive axillary lymph nodes, for disease-free survival.

PubMed ID: 24934787

Citation: Shulman LN, Berry DA, Cirrincione CT, Becker HP, Perez EA, O'Regan R, Martino S, Shapiro CL, Schneider CJ, Kimmick G, Burstein HJ, Norton L, Muss H, Hudis CA, Winer EP. Comparison of doxorubicin and cyclophosphamide versus single-agent paclitaxel as adjuvant therapy for breast cancer in women with 0 to 3 positive axillary nodes: CALGB 40101 (Alliance). J Clin Oncol. 2014 Aug 1;32(22):2311-7.

Associated Datasets: NCT00041119_D3_(FactorB_Patients)
NCT00041119_D4_(FactorB_AE)

Dataset Information

Dataset Name: NCT00041119_D4_(FactorB_AE)

Description: The NCT00041119_D4_(FactorB_AE) dataset is one of 2 datasets associated with PubMed ID 24934787. This dataset contains data on toxicity reported during treatment.

NCT00041119 D4 (FactorB AE) Data Dictionary

Variable description	Variable name	Codes	Notes
De-identified ID	MASK_ID		patient ID
Adeers report submitted	AER_SUBMITTED	1=No 2=Yes	
Meddra code	MEDDRA_CODE		
Meddra dictionary version	MEDDRA_VERSION		
AE grade	GRADE_ID		Grades 3 or 4 adverse events included
Event category	Eventcat		
Event description	Eventname		
AE related	RELATION_ID	1=Unrelated 2=Unlikely 3=Possible 4=Probable 5=Definitely	